arXiv:math/0412192v2 [math.QA] 10 Jan 2005

To L.D. Faddeev
with the respect and appreciation

Cayley-Hamilton theorem for quantum matrix algebras of $GL(m|n)$ type

D.I. Gurevich*
ISTV, Université de Valenciennes, 59304 Valenciennes, France
P.N. Pyatov†
Bogoliubov Laboratory of Theoretical Physics, JINR, 141980 Dubna, Russia
P.A. Saponov‡
Department of Theoretical Physics, IHEP, 142281 Protvino, Russia

February 1, 2008

Abstract

The classical Cayley-Hamilton identities are generalized to quantum matrix algebras of the $GL(m|n)$ type.

1 Introduction

The well known Cayley-Hamilton theorem of classical matrix analysis claims that any square matrix M with entries from a field $\mathbb{K}$ satisfies the identity

$$\Delta(M) \equiv 0, \tag{1.1}$$

where $\Delta(x) = \det(M - xI)$ is the characteristic polynomial of M. Provided $\mathbb{K}$ be algebraically closed, the coefficients of this polynomial are the elementary symmetric functions in the

*Dimitri.Gourevitch@univ-valenciennes.fr
†pyatov@thsun1.jinr.ru
‡Pavel.Saponov@ihep.ru

eigenvalues of M. Below for the sake of simplicity we assume $\mathbb{K}$ to be the field $\mathbb{C}$ of complex numbers.

In [NT, PS, GPS, IOPS, IOP1] the Cayley-Hamilton theorem was successively generalized to the case of the quantum matrix algebras of the $GL(m)$ type. Any such an algebra is constructed by means of R-matrix representations of Hecke algebras of the $GL(n)$ type.

In the works [KT] the Cayley-Hamilton identity was established for matrix superalgebras. The aim of the present paper consists in a further generalization of the results of [KT] to the case of quantum matrix superalgebras.

Let us formulate our main result — the Cayley-Hamilton theorem for the quantum matrix algebras of the $GL(m|n)$ type. All the definitions and notations are explained in the main text below.

Theorem. *Let $M = \|M_i^j\|_{i,j=1}^N$ be the matrix of the generators of the quantum matrix algebra $\mathcal{M}(R,F)$ defined by a pair of compatible, strictly skew-invertible R-matrices R and F, where R is a $GL(m|n)$ type R-matrix (see sections 3.1 and 3.2). Then in $\mathcal{M}(R,F)$ the following matrix identity holds*

$$\sum_{i=0}^{n+m} M^{\overline{m+n-i}}\, C_i \equiv 0\,, \tag{1.2}$$

where $M^{\bar{i}}$ is the i-th power of the quantum matrix (see definition (3.27)) and the coefficients C_i are linear combinations of the Schur functions (see definition (3.21)).

The Schur functions entering the expression of C_i are homogeneous polynomials of the $(mn+i)$ order in the generators M_i^j. Therefore, any matrix element in the left hand side of (1.2) is a homogeneous polynomial of the $(mn+m+n)$ order in M_i^j.

The expression of the coefficient C_i in terms of the Schur functions can be graphically presented as follows

$$C_i = \sum_{k=\max\{0,(i-n)\}}^{\min\{i,m\}} (-1)^k q^{(2k-i)} \tag{1.3}$$



where each Young diagram stands for the corresponding Schur function.

If the matrices F and R are the transpositions in the superspace $(m|n)$, then the corresponding algebra $\mathcal{M}(R,F)$ is a matrix superalgebra. In this case identity (1.2) coincides with the invariant Cayley-Hamilton relations obtained in [KT]. Note that in the supersymmetric case the linear size N of the matrix M is connected with the parameters m and n by the equation $N = m+n$. As was shown in [G], in general this relation does not take place in the quantum case. Our proof of identity (1.2) does not rely on such like connections.

The structure of the work is as follows. Section 2 contains definitions and necessary facts on the Hecke algebras. In section 3 we define the quantum matrix algebra of the $GL(m|n)$ type. Ibidem the notions of the characteristic subalgebra and of the matrix power space are introduced. The last section is devoted to the proof of the Cayley-Hamilton identity (1.2).

The authors express their appreciation to Alexei Davydov, Dimitry Leites, Alexander Molev and Hovhannes Khudaverdian. Special gratitude to our permanent co-authors Alexei Isaev and Oleg Ogievetsky for numerous fruitful discussions and remarks. The work of the second author (P.N.P.) was partially supported by the RFBR grant No. 03-01-00781.

2 Some facts on Hecke algebras

In this section we give a summary of facts on the structure of the Hecke algebras of A_{n-1} series. For their proofs the reader is referred to the works [DJ, M]. Detailed consideration of the material and the list of references on the theme can also be found in the reviews [OP, R]. When considering the quantum matrix algebras we shall use this section as a base of our technical tools.

2.1 Young diagrams and tableaux

First of all we recall some definitions from the partition theory. Mainly we shall use the terminology and notations of the book [Mac].

A partition λ of a positive integer n (notation: $\lambda \vdash n$) is a non-increasing sequence of non-negative integers λ_i such that their sum is equal to n:

$$\lambda = (\lambda_1, \lambda_2, \ldots), \quad \lambda_i \geq \lambda_{i+1} : \quad |\lambda| \equiv \sum \lambda_i = n.$$

The number $|\lambda| = n$ is called *the weight* of the partition λ. A convenient graphical image for a partition is *a Young diagram* (see [Mac]) and below we shall identify any partition with the corresponding Young diagram.

With each node of a Young diagram located at the intersection of the s-th column and the r-th row we associate a number $c = s - r$ which will be called *the content* of the node.

Define an inclusion relation $\subset$ on the set of all the Young diagrams by stating that a diagram μ includes a diagram λ (symbolically $\lambda \subset \mu$) if after the superposition the diagram λ is entirely contained inside of μ. In other words, $\lambda \subset \mu \Leftrightarrow \lambda_i \leq \mu_i \; i = 1, 2, \ldots$

Given a Young diagram $\lambda \vdash n$ one can construct $n!$ *Young tableaux* by writing into the nodes of the diagram all integers from 1 to n in an arbitrary order. The tableau will be noted by the symbol

$$\left\{ {\lambda \atop \alpha} \right\}, \qquad \alpha = 1, 2, \ldots, n!, \tag{2.1}$$

where the index α marks the different tableaux. The diagram λ will be called *the form* of a tableau $\left\{ {\lambda \atop \alpha} \right\}$.

The set of all tableaux associated with a diagram $\lambda \vdash n$ can be naturally supplied with an action of the n-th order symmetric group S_n. By definition, an element $\pi \in S_n$ acts on a tableaux $\left\{ {\lambda \atop \alpha} \right\}$ by changing any number i for $\pi(i)$ in all the nodes of the tableau. The resulting tableau will be denoted as $\left\{ {\lambda \atop \pi(\alpha)} \right\}$.

In the set of all the Young tableaux of the form λ we shall distinguish a subset of *standard* tableaux which satisfy an additional requirement: the numbers written in a standard tableau increases in any column from top to bottom and in any raw from left to right. From now on we shall use only standard Young tableaux and retain notation (2.1) for them. However, the index α will vary from 1 to an integer d_λ which is equal to the total number of all the standard Young tableaux of the same form λ. Explicit expressions for d_λ are given, for example, in [Mac].

Though the subset of the standard tableaux is not closed with respect to the action of the symmetric group, it nevertheless is a cyclic set under such an action since an arbitrary standard Young tableau of the form $\lambda \vdash n$ can be obtained from any other standard tableau of the same form by the action of some element π from the group S_n that is

$$\forall\, \alpha, \beta \quad \exists\, \pi \in S_n : \quad \left\{ {\lambda \atop \alpha} \right\} \mapsto \left\{ {\lambda \atop \beta} \right\} \equiv \left\{ {\lambda \atop \pi(\alpha)} \right\}.$$

The relation $\subset$ defined above for the Young diagrams can be extended to the set of the standard tableaux. Namely, we say that a tableau includes another one

$$\left\{ {\lambda \atop \alpha} \right\} \subset \left\{ {\mu \atop \beta} \right\},$$

if $\lambda \subset \mu$ and the numbers from 1 to $|\lambda|$ occupy the same nodes in the tableau $\left\{ {\mu \atop \beta} \right\}$ as they do in $\left\{ {\lambda \atop \alpha} \right\}$.

2.2 Hecke algebras: definition and the basis of matrix units

The Hecke algebra $\mathcal{H}_n(q)$ of A_{n-1} series is a unital associative $\mathbb{C}$-algebra generated by the set of elements $\{\sigma_i\}_{i=1}^{n-1}$ subject to the relations

$$\sigma_k \sigma_{k+1} \sigma_k = \sigma_{k+1} \sigma_k \sigma_{k+1} \,, \qquad k = 1, \ldots, n-2 \,, \tag{2.2}$$

$$\sigma_k \sigma_j = \sigma_j \sigma_k \,, \qquad \text{if } \; |k-j| \geq 2 \,, \tag{2.3}$$

$${\sigma_k}^2 = \mathbf{1} + (q - q^{-1}) \sigma_k \,, \qquad k = 1, \ldots, n-1 \,, \tag{2.4}$$

where $\mathbf{1}$ is the unit element. The last of the above defining relations depends on a parameter $q \in \mathbb{C} \setminus \{0\}$ and is referred to as *the Hecke condition*.

$\mathcal{H}_n(1)$ is identical with the group algebra $\mathbb{C}[S_n]$ of the symmetric group S_n. In case

$$k_q := \frac{q^k - q^{-k}}{q - q^{-1}} \neq 0 \,, \quad \text{for all} \quad k = 2, 3, \ldots, n, \tag{2.5}$$

the algebra $\mathcal{H}_n(q)$ is semisimple and is isomorphic to $\mathbb{C}[S_n]$. In what follows we shall suppose condition (2.5) to be fulfilled for all $k \in \mathbb{N} \setminus \{0\}$. Under such an assumption the algebra $\mathcal{H}_n(q)$ is isomorphic to the direct sum of matrix algebras

$$\mathcal{H}_n(q) \cong \bigoplus_{\lambda \vdash n} \mathrm{Mat}_{d_\lambda}(\mathbb{C}). \tag{2.6}$$

Here $\mathrm{Mat}_m(\mathbb{C})$ stands for an associative $\mathbb{C}$-algebra generated by m^2 matrix units E_{ab} $1 \leq a, b \leq m$ which satisfy the following multiplication law

$$E_{ab}\, E_{cd} = \delta_{bc}\, E_{ad}.$$

Due to isomorphism (2.6) one can construct the set of elements $E^\lambda_{\alpha\beta} \in \mathcal{H}_n(q)$, $\forall\, \lambda \vdash n$, $1 \leq \alpha, \beta \leq d_\lambda$, which are the images of matrix units generating the term Mat_{d_λ} in the direct sum (2.6)

$$E^\lambda_{\alpha\beta}\, E^\mu_{\gamma\tau} = \delta^{\lambda\mu} \delta_{\beta\gamma}\, E^\lambda_{\alpha\tau}. \tag{2.7}$$

The diagonal matrix units $E^\lambda_{\alpha\alpha} \equiv E^\lambda_\alpha$ are the primitive idempotents of the algebra $\mathcal{H}_n(q)$ and the set of all $E^\lambda_{\alpha\beta}$ forms a linear basis in $\mathcal{H}_n(q)$.

Obviously, the matrix units $E^\lambda_{\alpha\beta}$ can be realized in many different ways. The construction presented in [M, R] (see also [OP]) satisfies a number of useful additional relations. It is this type of the basis of matrix units which we shall use below. Now we shortly outline its main properties.

Firstly, in this basis there are known remarkably simple expressions for the elements which transform different diagonal matrix units into each other. In order to describe these elements we need a few new definitions.

With each generator σ_i of the algebra $H_n(q)$ we associate the element

$$\sigma_i(x) := \sigma_i + \frac{q^{-x}}{x_q}\, \mathbf{1}, \quad 1 \leq i \leq n-1. \tag{2.8}$$

Here x is an arbitrary non-negative integer, the symbol x_q being defined in (2.5). The elements $\sigma_i(x)$ obey the relations

$$\sigma_i(x)\sigma_{i+1}(x+y)\sigma_i(y) = \sigma_{i+1}(y)\sigma_i(x+y)\sigma_{i+1}(x), \qquad 1 \leq i \leq n-2, \tag{2.9}$$

$$\sigma_i(x)\sigma_i(-x) = \frac{(x+1)_q (x-1)_q}{x_q^2}\, \mathbf{1}, \qquad 1 \leq i \leq n-1. \tag{2.10}$$

Consider an arbitrary standard Young tableau $\left\{ {\lambda \atop \alpha} \right\}$ corresponding to a partition $\lambda \vdash n$. For each integer $1 \leq k \leq n-1$ we define the number $\ell_k\{{\lambda \atop \alpha}\}$ to be equal to the difference of the contents of nodes containing the numbers k and $(k+1)$

$$\ell_k\{{\lambda \atop \alpha}\} := c(k) - c(k+1). \tag{2.11}$$

Let a tableau $\left\{{\lambda \atop \beta}\right\}$ (not obligatory standard) differs from the initial tableau $\left\{{\lambda \atop \alpha}\right\}$ only in the transposition of the nodes with the numbers k and $(k+1)$

$$\beta = \pi_k(\alpha), \quad 1 \leq k \leq n-1, \tag{2.12}$$

$\pi_k \in S_n$ being the transposition of k and $(k+1)$. If the tableau $\left\{{\lambda \atop \pi_k(\alpha)}\right\}$ is standard then the following relations hold

$$\sigma_k(\ell_k\{{\lambda \atop \alpha}\})\, E^\lambda_\alpha = E^\lambda_{\pi_k(\alpha)}\, \sigma_k(-\ell_k\{{\lambda \atop \alpha}\})\,, \tag{2.13}$$

$$E^\lambda_\alpha\, \sigma_k(\ell_k\{{\lambda \atop \alpha}\}) = \sigma_k(-\ell_k\{{\lambda \atop \alpha}\})\, E^\lambda_{\pi_k(\alpha)}\,. \tag{2.14}$$

Now, take into account the cyclic property of the set of standard Young tableaux under the action of the symmetric group and the fact that an arbitrary permutation can be (non-uniquely) expanded into a product of transpositions. This allows one to connect any pair E^λ_α and E^λ_β by a chain of transformations similar to (2.13), (2.14). Moreover, relations (2.9), (2.10) ensure the compatibility of all possible formulae expressing dependencies between E^λ_α and E^λ_β.

If the tableau $\left\{{\lambda \atop \pi_k(\alpha)}\right\}$ is non-standard then

$$\sigma_k(\ell_k\{{\lambda \atop \alpha}\})\, E^\lambda_\alpha = E^\lambda_\alpha\, \sigma_k(\ell_k\{{\lambda \atop \alpha}\}) = 0. \tag{2.15}$$

The expressions for non-diagonal matrix units follow from (2.13), (2.14). Assuming both the tableaux $\left\{{\lambda \atop \alpha}\right\}$ and $\left\{{\lambda \atop \pi_k(\alpha)}\right\}$ to be standard, one gets

$$E^\lambda_{\alpha\pi_k(\alpha)} := \omega(\ell_k\{{\lambda \atop \alpha}\})\, E^\lambda_\alpha\, \sigma_k(\ell_k\{{\lambda \atop \alpha}\})\,, \tag{2.16}$$

$$E^\lambda_{\pi_k(\alpha)\alpha} := \omega(\ell_k\{{\lambda \atop \alpha}\})\, \sigma_k(\ell_k\{{\lambda \atop \alpha}\})\, E^\lambda_\alpha\,, \tag{2.17}$$

where the normalizing coefficient $\omega(\ell)$ satisfies the relation

$$\omega(\ell)\, \omega(-\ell) = \frac{\ell_q^2}{(\ell+1)_q(\ell-1)_q}\,. \tag{2.18}$$

In particular, one can set $\omega(\ell) := \ell_q/(\ell_q+1)$. Note that in case the tableau $\left\{{\lambda \atop \pi_k(\alpha)}\right\}$ is a standard one, we have $\ell_k\{{\lambda \atop \alpha}\} \neq \pm 1$. Therefore, the above expression for the normalizing coefficient is always correctly defined.

To formulate the second property of the matrix units which will be used below let us consider the chain of Hecke algebra embeddings

$$\mathcal{H}_2(q) \hookrightarrow \ldots \hookrightarrow \mathcal{H}_k(q) \hookrightarrow \mathcal{H}_{k+1}(q) \hookrightarrow \ldots, \tag{2.19}$$

which are defined on generators by the formula

$$\mathcal{H}_k(q) \ni \sigma_i \mapsto \sigma_{i+1} \in \mathcal{H}_{k+1}(q), \quad i = 1, \ldots, k-1. \tag{2.20}$$

In each subalgebra $\mathcal{H}_k(q)$ one can choose the basis of the matrix units $E^{\lambda}_{\alpha\beta}$ in such a way that the diagonal matrix units $E^{\lambda}_{\alpha} \in \mathcal{H}_k(q)$ can be decomposed into the sum of diagonal matrix units belonging to any enveloping algebra $\mathcal{H}_m(q) \supset \mathcal{H}_k(q)$

$$E^{\lambda\vdash k}_{\alpha} = \sum_{\left\{{\lambda \atop \alpha}\right\} \subset \left\{{\mu \atop \beta}\right\}} E^{\mu\vdash m}_{\beta}, \quad m \geq k. \tag{2.21}$$

Here the summation goes over all the standard Young tableaux $\left\{{\mu \atop \beta}\right\}$, $\mu \vdash m$, containing the standard tableau $\left\{{\lambda \atop \alpha}\right\}$.

3 Quantum matrix algebra $\mathcal{M}(R, F)$

3.1 R-matrix representations of the Hecke algebra

Let V be a finite dimensional vector space over the field of complex numbers, $\dim V = N$. With any element $X \in \mathrm{End}(V^{\otimes p})$, $p = 1, 2, \ldots$, we associate a sequence of endomorphisms $X_i \in \mathrm{End}(V^{\otimes k})$, $k = p, p+1, \ldots$, according to the rule

$$X_i = \mathrm{Id}_V^{i-1} \otimes X \otimes \mathrm{Id}_V^{k-p-i+1}, \quad 1 \leq i \leq k-p+1, \tag{3.1}$$

where Id_V is the identical automorphism of V.

An operator $R \in \mathrm{Aut}(V^{\otimes 2})$ satisfying the *Yang-Baxter equation*

$$R_1 R_2 R_1 = R_2 R_1 R_2\,, \tag{3.2}$$

will be called *the R-matrix*. One can easily see that any R-matrix obeying the condition

$$(R - q\,\mathrm{Id}_{V^{\otimes 2}})(R + q^{-1}\,\mathrm{Id}_{V^{\otimes 2}}) = 0, \quad q \in \mathbb{C} \setminus \{0\} \tag{3.3}$$

generates representations ρ_R for the series of the Hecke algebras $\mathcal{H}_k(q)$, $k = 2, 3, \ldots$

$$\rho_R : \mathcal{H}_k(q) \to \mathrm{End}(V^{\otimes k}) \qquad \sigma_i \mapsto \rho_R(\sigma_i) = R_i \qquad 1 \leq i \leq k-1, \tag{3.4}$$

where σ_i are the generators of $\mathcal{H}_k(q)$ (2.2) – (2.4). Such like R-matrices will be called *the Hecke type* (or simply *Hecke*) R-matrices and the corresponding representations ρ_R — *the R-matrix* ones.

Let R be a Hecke R-matrix. Suppose that the corresponding representations $\rho_R : \mathcal{H}_k(q) \to \mathrm{End}(V^{\otimes k})$ are faithful for all $2 \leq k < (m+1)(n+1)$ while the representation $\rho_R : \mathcal{H}_{(m+1)(n+1)}(q) \to \mathrm{End}(V^{\otimes (m+1)(n+1)})$ possesses a kernel, generated by any matrix unit corresponding to the rectangular Young diagram $((n+1)^{(m+1)})$ that is

$$\begin{aligned} &\rho_R(E^{((n+1)^{(m+1)})}_{\alpha}) = 0 \\ &\rho_R(E^{\mu}_{\alpha}) \neq 0, \qquad \text{for any } \mu \vdash (m+1)(n+1), \quad \mu \neq ((n+1)^{(m+1)}). \end{aligned} \tag{3.5}$$

Such an R-matrix will be referred to as an R-matrix of *the $GL(m|n)$ type*.

3.2 R-trace and pairs of compatible R-matrices

Let $\{v_i\}_{i=1}^N$ be a fixed basis in the space V. In the basis $\{v_i \otimes v_j\}_{i,j=1}^N$ an operator $X \in \mathrm{End}(V^{\otimes 2})$ is defined by its matrix X_{ij}^{kl}: $X(v_i \otimes v_j) := \sum_{k,l=1}^N X_{ij}^{kl}\, v_k \otimes v_l$.

The operator $X \in \mathrm{End}(V^{\otimes 2})$ will be called *skew-invertible* if there exists a matrix ${\Psi^X}_{ij}^{kl}$ such that

$$\sum_{a,b=1}^{N} X_{ia}^{kb} {\Psi^X}_{bj}^{al} = \delta_i^l\, \delta_j^k. \tag{3.6}$$

One can easily see that the property of skew-invertibility is invariant with respect to the change of the basis $\{v_i\}$ and, therefore, to the matrix ${\Psi^X}_{ij}^{kl}$ there corresponds an operator $\Psi^X \in \mathrm{End}(\mathrm{V}^{\otimes 2})$. Introduce a notation

$$D^X := \mathrm{Tr}_{(2)} \Psi^X \in \mathrm{End}(V), \tag{3.7}$$

where the symbol $\mathrm{Tr}_{(2)}$ means taking the trace over the second space in the product $V \otimes V$. A skew-invertible operator X will be called *strictly* skew-invertible in case the operator D^X is invertible.

Consider the set of $N \times N$ matrices $\mathrm{Mat}_N(W)$, their entries being vectors of a linear space W over $\mathbb{C}$. The set $\mathrm{Mat}_N(W)$ is endowed with the structure of a linear space over $\mathbb{C}$ in the obvious way.

Let R be a skew-invertible R-matrix. The linear map

$$\mathrm{Tr}_R : \mathrm{Mat}_N(W) \to W,$$

defined by

$$\mathrm{Tr}_R(M) = \sum_{i,j=1}^{N} {D^R}_i^j M_j^i \qquad M \in \mathrm{Mat}_N(W)\,, \tag{3.8}$$

will be called the calculation of *R-trace*.

Remark 1 If as an R-matrix one chooses the transposition operator in the superspace of $(m|n)$ type (see definition (3.32)), then the operation Tr_R coincides with the calculation of the supertrace.

An ordered pair $\{R, F\}$ of the two R-matrices R and F will be called *a compatible pair* if the following relations are satisfied

$$\begin{aligned} R_1 F_2 F_1 &= F_2 F_1 R_2 \\ R_2 F_1 F_2 &= F_1 F_2 R_1. \end{aligned} \tag{3.9}$$

Now we list some properties of the compatible pairs of R-matrices.

1. If R is a skew-invertible R-matrix and $\{R, F\}$ is a compatible pair, then for all $M \in \mathrm{Mat}_N(W)$ the following relation holds

$$\mathrm{Tr}_{R^{(2)}}(F_1^{\pm 1} M_1 F_1^{\mp 1}) = \mathrm{Id}_V \, \mathrm{Tr}_R M \quad \text{where} \quad M_1 = M \otimes \mathrm{Id}_V. \tag{3.10}$$

2. In case R is skew-invertible, then

$$[R_1, D_1^R D_2^R] = 0\,. \tag{3.11}$$

   A direct consequence of this formula is a *cyclic property* of the R-trace

$$\mathrm{Tr}_{R^{(12)}}(R_1 U) = \mathrm{Tr}_{R^{(12)}}(U R_1), \quad U \in \mathrm{Mat}_N(W)^{\otimes 2}. \tag{3.12}$$

3. If $\{R, F\}$ is a compatible pair, then $R_f := F^{-1} R^{-1} F$ is an R-matrix and the pair $\{R_f, F\}$ is compatible.

4. Let $\{R, F\}$ be a compatible pair, R-matrix R_f is skew-invertible and matrices R and F are strictly skew-invertible. Then the map $\phi : \mathrm{Mat}_N(W) \to \mathrm{Mat}_N(W)$ defined by

$$\phi(M) := \mathrm{Tr}_{R^{(2)}}(F_1 M_1 F_1^{-1} R_1), \quad M \in \mathrm{Mat}_N(W), \tag{3.13}$$

   is invertible. An explicit form of its inverse ϕ^{-1} is as follows (see [OP2])

$$\phi^{-1}(M) = \mathrm{Tr}_{R_f{}^{(2)}}(F_1^{-1} M_1 R_1^{-1} F_1). \tag{3.14}$$

3.3 Quantum matrix algebras: definition

Consider a linear space $\mathrm{Mat}_N(W)$ and define a series of linear mappings $\mathrm{Mat}_N(W)^{\otimes k} \to \mathrm{Mat}_N(W)^{\otimes (k+1)}$, $k \geq 1$, by the following recurrent relations

$$M_{\overline{1}} := M, \quad M_{\overline{k}} \mapsto M_{\overline{k+1}} := F_k M_{\overline{k}} F_k^{-1}, \quad M_{\overline{k}} \in \mathrm{Mat}_N(W)^{\otimes k}, \tag{3.15}$$

where M is an arbitrary matrix from $\mathrm{Mat}_N(W)$ and F is a fixed element of $\mathrm{Aut}(V \otimes V)$.

As the linear space W we choose an associative algebra $\mathcal{A}$ over $\mathbb{C}$ freely generated by the unit element and N^2 generators M_i^j

$$\mathcal{A} = \mathbb{C}\langle M_i^j \rangle \quad 1 \leq i, j \leq N.$$

Definition 2 *Let $\{R, F\}$ be a compatible pair of strictly skew-invertible R-matrices, the R-matrix $R_f = F^{-1} R^{-1} F$ being skew-invertible. By definition, the quantum matrix algebra $\mathcal{M}(R, F)$ is a quotient of the algebra $\mathcal{A}$ over a two-sided ideal generated by entries of the following matrix relation*

$$R_1 M_{\overline{1}} M_{\overline{2}} - M_{\overline{1}} M_{\overline{2}} R_1 = 0, \tag{3.16}$$

where $M = \|M_i^j\| \in \mathrm{Mat}_N(\mathcal{A})$ and the matrices $M_{\overline{k}}$ are constructed according to (3.15) with the R-matrix F.

Remark 3 The definition of the quantum matrix algebra presented here differs from that given in [IOP1]. Nevertheless, under the additional assumptions on R and F made above, both definitions are equivalent. As was shown in [IOP2], the matrix $\widetilde{M} = (D^F)^{-1} M D^R$ obeys relations imposed on the generators of the quantum matrix algebra in the work [IOP1].

Lemma 4 *The matrix M composed of the generators of the quantum matrix algebra $\mathcal{M}(R,F)$ satisfies the relations*

$$R_k\, M_{\overline{k}}\, M_{\overline{k+1}} = M_{\overline{k}}\, M_{\overline{k+1}}\, R_k, \tag{3.17}$$

where the matrices $M_{\overline{k}}$, $M_{\overline{k+1}}$ are defined by rules (3.15) with the R-matrix F.

The proof of the lemma is presented in [IOP1].

3.4 Characteristic subalgebra and Schur functions

Further in this section we consider the quantum matrix algebras $\mathcal{M}(R,F)$ defined by a Hecke R-matrix R. We shall refer to them as the matrix algebras of *the Hecke type*.

Let a subspace $\mathrm{Char}(R,F) \subset \mathcal{M}(R,F)$ be a linear span of the unit and of the following elements

$$y(x^{(k)}) = \mathrm{Tr}_{R^{(1\,\ldots\,k)}}(M_{\overline{1}} \ldots M_{\overline{k}}\, \rho_R(x^{(k)})) \quad k = 1, 2, \ldots, \tag{3.18}$$

where $x^{(k)}$ runs over all elements of $\mathcal{H}_k(q)$. The symbol $\mathrm{Tr}_{R^{(1\,\ldots\,k)}}$ stands for the calculation of the R-trace over the spaces from the first to the k-th ones.

Proposition 5 *Let $\mathcal{M}(R,F)$ be a quantum matrix algebra of the Hecke type. The subspace $\mathrm{Char}(R,F)$ is a commutative subalgebra of $\mathcal{M}(R,F)$. The subalgebra $\mathrm{Char}(R,F)$ will be called the characteristic subalgebra.*

The detailed proof is given in [IOP1]. It is based, in particular, on the following technical result.

Lemma 6 *Consider an arbitrary element $x^{(k)}$ of the algebra $\mathcal{H}_k(q)$. Let $x^{(k)\uparrow i}$ denotes the image of $x^{(k)}$ in the algebra $\mathcal{H}_{k+i}(q)$ under the embedding $\mathcal{H}_k(q) \hookrightarrow \mathcal{H}_{k+i}(q)$ defined by (2.20). Let $\{R,F\}$ be a compatible pair of R-matrices, the Hecke R-matrix R being skew-invertible. Then the following relation holds true*

$$\mathrm{Tr}_{R^{(i+1,\ldots,i+k)}}(M_{\overline{i+1}} \ldots M_{\overline{i+k}}\; \rho_R(x^{(k)\uparrow i})) \;=\; \mathrm{Id}_{V^{\otimes i}}\; y(x^{(k)}). \tag{3.19}$$

Formula (3.19) will be repeatedly used in the sequel. It is immediate consequence of the property (3.10) of the compatible pairs $\{R,F\}$ and its proof is also presented in [IOP1].

Take into consideration two sets $\{p_k(M)\}$ and $\{s_\lambda(M)\}$ of elements of the characteristic subalgebra. The elements of these sets will be referred to as *the power sums* and *the Schur functions* respectively. For an arbitrary integer $k \geq 1$ the power sums are defined by

$$p_k(M) := \mathrm{Tr}_{R^{(1\,\ldots\,k)}}(M_{\overline{1}} \ldots M_{\overline{k}}\, R_{k-1} \ldots R_1). \tag{3.20}$$

Given a Hecke type R-matrix R and an arbitrary partition $\lambda \vdash k$, $k = 1, 2, \ldots$, we define the Schur functions as follows

$$s_0(M) := 1, \quad s_\lambda(M) := \mathrm{Tr}_{R^{(1\ldots k)}}(M_{\overline{1}} \ldots M_{\overline{k}}\, \rho_R(E^\lambda_\alpha)). \tag{3.21}$$

The right hand side of (3.21) does not actually depend on the index α of the matrix unit. Indeed, for all $x^{(k)} \in \mathcal{H}_k(q)$ the matrices $\rho_R(x^{(k)})$ and $(M_{\overline{1}} \ldots M_{\overline{k}})$ commute in (3.18) due to (3.17). Then, taking into account the cyclic property of the R-trace, one gets

$$y\left(\sigma_k(\ell)\, E^\lambda_\alpha\, \sigma_k^{-1}(\ell)\right) \;=\; y(E^\lambda_\alpha)\,, \quad y(E^\lambda_{\pi_k(\alpha)\alpha}) = w(\ell)\, y\left(E^\lambda_{\pi_k(\alpha)}\, \sigma_k(\ell)\, E^\lambda_\alpha\right) = 0, \tag{3.22}$$

where $\ell := \ell_k\{{}^\lambda_\alpha\}$. Next, with the use of (2.8), (2.10), (2.17) and (2.18) relations (2.13) can be rewritten in the form

$$E^\lambda_{\pi_k(\alpha)} \;=\; \sigma_k(\ell)\, E^\lambda_\alpha\, \sigma_k^{-1}(\ell) \;-\; (q - q^{-1})\, w(-\ell)\, E^\lambda_{\pi_k(\alpha)\alpha}\,. \tag{3.23}$$

This leads to $y(E^\lambda_{\pi_k(\alpha)}) = y(E^\lambda_\alpha)$. At last, the fact that the set of the standard Young tableaux is cyclic with respect to the action of the symmetric group guarantees that definition (3.21) is non-contradictory.

Proposition 7 *For any quantum matrix algebra $\mathcal{M}(R,F)$ of the Hecke type*

i) Its characteristic subalgebra $\mathrm{Char}(R, F)$ *is generated by the unit and by the power sums $p_k(M)$, $k = 1, 2, \ldots$;*

ii) The set of Schur functions $s_\lambda(M)$, $\lambda \vdash k$, $k = 0, 1, \ldots$, forms a linear basis in $\mathrm{Char}(R, F)$.

Proof. The first part of the proposition is proved in [IOP1]. To prove the second part, we note that, basing on the arguments given to justify relations (3.21), one can get the following equalities

$$y(E^\lambda_{\alpha\beta}) \;=\; \delta_{\alpha\beta}\, y(E^\lambda_\alpha), \quad y(E^\lambda_\alpha) \;=\; y(E^\lambda_\beta), \qquad \forall\ \alpha, \beta. \tag{3.24}$$

Now the statement on a linear basis of the Schur functions follows from the fact that the matrix units $\{E^\lambda_{\alpha\beta} \,|\, \lambda \vdash k\}$ form the linear basis in the algebra $\mathcal{H}_k(q)$. □

3.5 Powers of quantum matrices

Let a subspace $\mathrm{Pow}(R, F) \subset \mathrm{Mat}_N(\mathcal{M}(R, F))$ be the linear span of the unit matrix and of the following matrices

$$M^{(x^{(k)})} := \mathrm{Tr}_{R^{(2\ldots k)}}(M_{\overline{1}} \ldots M_{\overline{k}}\, \rho_R(x^{(k)})) \qquad k = 1, 2, \ldots, \tag{3.25}$$

where $x^{(k)}$ runs over all elements of $\mathcal{H}_k(q)$. The matrix $M^{(x^{(k)})}$ will be called *the $x^{(k)}$-th power* of the matrix M, and the space $\mathrm{Pow}(R, F)$ will be referred to as *the space of matrix powers*.

Proposition 8 $\mathrm{Pow}(R,F)$ *is a right module over the characteristic subalgebra* $\mathrm{Char}(R,F)$.

With the use of (3.19) this proposition can be proved analogously to the proof of the commutativity of $\mathrm{Char}(R,F)$ (see [IOP1]).

In the space $\mathrm{Pow}(R,F)$ we choose a special set of matrices constructed by the following rule. Having fixed a standard Young tableau $\left\{\begin{smallmatrix}\lambda\\ \alpha\end{smallmatrix}\right\}$ of the form $\lambda \vdash k$, $k = 1, 2, \ldots$, we define the matrices

$$M^{((1);1)} := M, \quad M^{(\lambda;i)} := \mathrm{Tr}_{R^{(2\ldots k)}}(M_{\overline{1}} \ldots M_{\overline{k}}\, \rho_R(E^{\lambda}_{\alpha})), \quad k = 2, 3, \ldots, \tag{3.26}$$

where the index i in $M^{(\lambda;i)}$ is the number of the raw of the tableau $\left\{\begin{smallmatrix}\lambda\\ \alpha\end{smallmatrix}\right\}$ (when counting from the top) which contains k — the largest integer among those filling the tableau. Take, as an example, the standard tableau

1	3	4	6
2	7		
5			

.

Since the largest integer $k = 7$ is contained in the second raw then $i = 2$ for the diagram involved.

The matrix $M^{(\lambda;i)}$ defined in (3.26) does not depend on the positions of the other integers which are less than k. The proof is analogous to that used when arguing in favour of the unambiguity of the Schur function definition (see the paragraph next to the formula (3.21)). One should only take into account an additional circumstance. If two matrix units E^{λ}_{α} and E^{λ}_{β} $\lambda \vdash k$ have the number k standing in the same nodes of the corresponding tableaux, then the chain of transformations linking these matrix units contains only the elements $x^{(k-1)}$ of the subalgebra $\mathcal{H}_{k-1}(q)$. Just with respect to the images $\rho_R(x^{(k-1)}) \in \mathrm{Id}_V \otimes \mathrm{Aut}(V^{\otimes(k-1)})$ the operation $\mathrm{Tr}_{R^{(2\ldots k)}}$ possesses the cyclic property.

Besides $M^{(\lambda;i)}$, consider also a series of matrices $M^{\overline{k}}$:

$$M^{\overline{1}} := M, \quad M^{\overline{k}} := \mathrm{Tr}_{R^{(2\ldots k)}}(M_{\overline{1}} \ldots M_{\overline{k}}\, R_{k-1} \ldots R_1), \quad k = 2, 3, \ldots. \tag{3.27}$$

The matrices $M^{(\lambda;i)}$ and $M^{\overline{k}}$ will be called respectively *the* $(\lambda;i)$*-th* and *the* k*-th powers* of the matrix M.

Proposition 9 *For any quantum matrix algebra* $\mathcal{M}(R,F)$ *of the Hecke type:*

i) The unit matrix and the family of matrices $M^{\overline{k}}$, $k = 1, 2, \ldots$ *is a generating set of the right* $\mathrm{Char}(R,F)$*-module* $\mathrm{Pow}(R,F)$*;*

ii) The unit matrix and the family of matrices $M^{(\lambda;i)}$, $\lambda \vdash k$, $k = 1, 2, \ldots$ *where* i *runs over all the possible values for each partition* $\lambda \vdash k$ *is a basis of the linear space* $\mathrm{Pow}(R,F)$.

The proof of items *i)* and *ii)* of the above proposition can be given by a quite obvious generalization of the proof of the corresponding items of proposition 7.

To end the section, we present another form of the k-th power of the matrix M. Basing on the property (3.10) of compatible pairs one can show that definition (3.27) is equivalent to the following iterative formulae

$$M^{\overline{0}} = \mathrm{Id}_V, \quad M^{\overline{k}} = M \cdot \phi(M^{\overline{k-1}}), \tag{3.28}$$

where the map ϕ is given by (3.13) and the dot stands for the usual matrix product. Relations (3.28) can be interpreted as the result of the consecutive application of the operator $\mathsf{M} : \mathrm{Mat}_N(\mathcal{M}(R,F)) \to \mathrm{Mat}_N(\mathcal{M}(R,F))$

$$\mathsf{M}(X) := M \cdot \phi(X), \quad X \in \mathrm{Mat}_N(\mathcal{M}(R,F))$$

to the unit matrix Id_V. Indeed, since $\phi(\mathrm{Id}_V) = \mathrm{Id}_V$ one gets

$$\mathsf{M}(\mathrm{Id}_V) = M, \quad M^{\overline{k}} = \mathsf{M}(M^{\overline{k-1}}) = \ldots = \mathsf{M}^k(\mathrm{Id}_V).$$

3.6 Quantum matrix algebras: examples

Let $P : V^{\otimes 2} \to V^{\otimes 2}$ be the transposition operator:

$$P(v_1 \otimes v_2) = v_2 \otimes v_1$$

and R-matrix R be strictly skew-invertible. The pair $\{R, P\}$ is compatible and it defines a quantum matrix algebra $\mathcal{M}(R,P)$. Denote the matrix of its generators by the letter T. In the case involved, $T_{\overline{k}} = T_k$ (see definitions (3.1) and (3.15)) and commutation relations (3.16) take the form

$$R_1 T_1 T_2 - T_1 T_2 R_1 = 0. \tag{3.29}$$

Suppose additionally that R is of the $GL(m)$ type. The standard example of such an R-matrix is the Drinfeld-Jimbo R-matrix obtained by the quantization of classical groups of the A_m series. The corresponding algebra $\mathcal{M}(R,P) = \mathrm{Fun}_q(GL(N))$ is the quantization of the algebra of functions on the general linear group [FRT]. Note that in general $m \neq N$. Examples of $N^2 \times N^2$ R-matrices of the $GL(m)$ type such that $m \neq N$ are constructed in [G].

In case R is an R-matrix of the $GL(m)$ type (including the Drinfeld-Jimbo R-matrix) the matrix T of generators of $\mathcal{M}(R,P)$ obeys a polynomial Cayley-Hamilton identity of the m-th order [IOPS, IOP1]. The coefficients of the polynomial are proportional to the Schur functions $s_{(1^k)}(T)$, $0 \leq k \leq m$. At the classical limit $q \to 1$ the Drinfeld-Jimbo R-matrix tends to the transposition matrix $\lim_{q\to 1} R = P$ and algebra (3.29) becomes the commutative matrix algebra $\mathcal{M}(P,P)$. The corresponding Cayley-Hamilton identity coincides with (1.1).

In particular, the Schur functions $s_{(1^k)}(T)$ turn into the elementary symmetric functions in the eigenvalues of the matrix T.

The other example of a compatible pair is given by $\{R, R\}$ where R is a strictly skew-invertible R-matrix. The commutation relations (3.16) for the generators of the quantum matrix algebra $\mathcal{M}(R, R)$ read

$$R_1 L_1 R_1 L_1 - L_1 R_1 L_1 R_1 = 0, \tag{3.30}$$

where $L_1 = L \otimes \mathrm{Id}_V$ and $L = \|L_i^j\|$ is the matrix composed of the algebra generators. The algebra $\mathcal{M}(R, R)$ is called *the reflection equation algebra* [KS]. Having first appeared in the theory of integrable systems with boundaries, the algebra has then found applications in the differential geometry of the quantum groups (see, for example, [IP, FP, GS, GS2]).

In this case the power of the quantum matrix coincides with the usual matrix power $L^{\overline{k}} = L^k$. The characteristic subalgebra $\mathrm{Char}(R, R)$ belongs to the center of $\mathcal{M}(R, R)$. The Cayley-Hamilton identity for the reflection equation algebra with Drinfeld-Jimbo R-matrix was obtained in [NT, PS]. Its generalization to the case of an arbitrary R-matrix of the $GL(m)$ type was found in [GPS].

Supposing R to be a strictly skew-invertible R-matrix of the Hecke type, let us make a linear shift of generators of $\mathcal{M}(R, R)$

$$L \mapsto \mathcal{L} = L + \frac{1}{q - q^{-1}} \mathrm{Id}_V.$$

After such a change relations (3.30) transforms into the quadratic-linear form

$$R_1 \mathcal{L}_1 R_1 \mathcal{L}_1 - \mathcal{L}_1 R_1 \mathcal{L}_1 R_1 = R_1 \mathcal{L}_1 - \mathcal{L}_1 R_1. \tag{3.31}$$

The above basis of generators is convenient in constructing the quantum analogs of orbits of the coadjoint action of a Lie group on the space dual to its Lie algebra. The Cayley-Hamilton identity for this basis was obtained in [GS]. If R is the Drinfeld-Jimbo R-matrix, then in the limit $q \to 1$ commutation relations (3.31) of the $\mathcal{M}(R, R)$ generators pass to that of the generators of the universal enveloping algebra $U(gl_N)$ and the Cayley-Hamilton identity transforms into the well known identity for the matrix composed of the $U(gl_N)$ generators (see, for example, [Gou]).

Let $P_{m|n}$ be the transposition operator on the superspace of the $(m|n)$ type

$$P_{m|n}(v_1 \otimes v_2) = (-1)^{|v_1||v_2|} v_2 \otimes v_1, \tag{3.32}$$

where v_i is a homogeneous element of the superspace and $|v_i|$ denotes the corresponding parity. The algebra $\mathcal{M}(R, R)$ with $R = P_{m|n}$ was considered in [KT]. Note that the matrix superalgebra $\mathcal{M}(P_{m|n}, P_{m|n})$ is a limiting case of the algebra $\mathcal{M}(R_{m|n}, R_{m|n})$ (3.30) at $q \to 1$ where $R_{m|n}$ is the Drinfeld-Jimbo R-matrix obtained in quantization of the classical supergroup $GL(m|n)$ (an explicit form of this R-matrix can be found in [DKS, I]). The so-called invariant Cayley-Hamilton identity established in [KT] is a limiting case of the Cayley-Hamilton identity for the quantum matrix algebras of the $GL(m|n)$ type. The latter is proved in the next section.

4 Proof of the Cayley-Hamilton identities

In this section we give a scheme of the proof and then formulate the Cayley-Hamilton identity for the quantum matrix algebras $\mathcal{M}(R,F)$ defined by an R-matrix R of the $GL(m|n)$ type. Below these algebras will be referred to as *the matrix algebras of the $GL(m|n)$ type*.

Fix a pair of integers $m \geq 1,\ n \geq 1$ and denote $A := (m+1)(n+1)$. We also introduce the special notations for the following partitions

$$\Lambda(r,s) := \Big((n+1)^r, n^{(m-r)}, s\Big), \quad r=0,\dots,m,\ s=0,\dots,n, \tag{4.1}$$

$$\Lambda^+(r,s) := \Big(n+2, (n+1)^{(r-1)}, n^{(m-r)}, s\Big), \quad r=1,\dots,m,\ s=0,\dots,n, \tag{4.2}$$

$$\Lambda_+(r,s) := \Big((n+1)^r, n^{(m-r)}, s, 1\Big), \quad r=0,\dots,m,\ s=1,\dots,n. \tag{4.3}$$

Depicted below are the corresponding Young diagrams

$\Lambda(r,s) =$ 

$\Lambda^+(r,s) =$ 

$\Lambda_+(r,s) =$ 

Let the symbol $E_{row}^{\Lambda(r,s)}$, $s \geq 1$, (respectively, $E_{col}^{\Lambda(r,s)}$, $r \geq 1$, $E^{\Lambda^+(r,s)}$, $E^{\Lambda_+(r,s)}$) denotes a diagonal matrix unit which corresponds to a standard tableau of the form $\Lambda(r,s)$ (respectively, $\Lambda(r,s)$, $\Lambda^+(r,s)$, $\Lambda_+(r,s)$) containing a standard Young tableau of the form $\Lambda(r,s-1)$ (respectively, $\Lambda(r-1,s)$, $\Lambda(r,s)$, $\Lambda(r,s)$). In other words, in the diagonal matrix units $E_{row}^{\Lambda(r,s)}$, $E_{col}^{\Lambda(r,s)}$, $E^{\Lambda^+(r,s)}$ and $E^{\Lambda_+(r,s)}$ one fixes the position of the largest number S (S is equal to $mn+r+s$, $mn+r+s$, $mn+r+s+1$ and $mn+r+s+1$ respectively) as is shown in the picture below

$E_{row}^{\Lambda(r,s)} =$ 

$E_{col}^{\Lambda(r,s)} =$ 

$E^{\Lambda^+(r,s)} =$ 

$E^{\Lambda_+(r,s)} =$ 

. (4.4)

Emphasize, that the further calculations do not depend on the positions of the other numbers in the nodes of the matrix units (4.4). At last, introduce the special notations for the following elements of the space $\mathrm{Mat}_N(\mathcal{M}(R,F))$

$$P_{row}(r,s) := \mathrm{Tr}_{R^{(2\dots A)}}\Big(M_{\overline{2}}\dots M_{\overline{A}}\ \rho_R(E_{row}^{\Lambda(r,s)})\ R_t R_{t-1}\dots R_1\Big), \tag{4.5}$$

$$P_{col}(r,s) := \mathrm{Tr}_{R^{(2\dots A)}}\Big(M_{\overline{2}}\dots M_{\overline{A}}\ \rho_R(E_{col}^{\Lambda(r,s)})\ R_t R_{t-1}\dots R_1\Big), \tag{4.6}$$

$$P^{+}(r,s) \;:=\; \mathrm{Tr}_{R^{(2\dots A)}}\left(M_{\overline{2}}\dots M_{\overline{A}}\;\rho_R(E^{\Lambda^{+}(r,s)})\,R_{t-1}R_{t-2}\dots R_1\right), \tag{4.7}$$

$$P_{+}(r,s) \;:=\; \mathrm{Tr}_{R^{(2\dots A)}}\left(M_{\overline{2}}\dots M_{\overline{A}}\;\rho_R(E^{\Lambda_{+}(r,s)})\,R_{t-1}R_{t-2}\dots R_1\right), \tag{4.8}$$

where $t := (m-r)+(n-s)+1$.

Let us take the following linear combinations of the above matrices

$$\begin{aligned}\Phi_i \;:=\; &\sum_{k=\max\{0,i-n\}}^{\min\{i-1,m\}}(-1)^k\,\frac{(i-k)_q(m+n-i+k+2)_q}{(m+n-i+2)_q}\,P_{row}(k,i-k)\\ &-\sum_{k=\max\{1,i-n\}}^{\min\{i,m\}}(-1)^k\,\frac{k_q(m+n-k+2)_q}{(m+n-i+2)_q}\,P_{col}(k,i-k)\;,\end{aligned} \tag{4.9}$$

where the index i runs from 1 to $(m+n)$. Our immediate task is in proving the relations

$$\Phi_{i+1}-\Phi_i \;=\; \phi(M^{\overline{m+n-i}})\sum_{k=\max\{0,i-n\}}^{\min\{i,m\}}(-1)^k\,q^{2k-i}\,s_{\Lambda(k,i-k)}(M)\,,\quad i=1,2,\dots,m+n-1. \tag{4.10}$$

With that end in view, transform the expression of Φ_i to the form

$$\begin{aligned}\Phi_i \;=\; &\sum_{k=\max\{0,i-n\}}^{\min\{i-1,m\}}(-1)^k\,\tfrac{(i-k)_q(m+n-i+k+2)_q}{(m+n-i+2)_q}\Big\{\tfrac{q^{(m+n+k-i+2)}}{(m+n+k-i+2)_q}P^{+}(k,i-k)\\ &-\tfrac{q^{-(i-k)}}{(i-k)_q}P_{+}(k,i-k)+qP_{row}(k,i-k+1)+\tfrac{q^{(m+n-i+1)}}{(m+n-i+1)_q}P_{col}(k+1,i-k)\Big\}\\ -\;&\sum_{k=\max\{1,i-n\}}^{\min\{i,m\}}(-1)^k\,\tfrac{k_q(m+n-k+2)_q}{(m+n-i+2)_q}\Big\{\tfrac{q^k}{k_q}P^{+}(k,i-k)-\tfrac{q^{-(m+n-k+2)}}{(m+n-k+2)_q}P_{+}(k,i-k)\\ &-\tfrac{q^{-(m+n-i+1)}}{(m+n-i+1)_q}P_{row}(k,i-k+1)-q^{-1}P_{col}(k+1,i-k)\Big\}.\end{aligned} \tag{4.11}$$

Here we extend the definition of elements (4.5)-(4.8) to the boundary values of its indices, that is to the cases when the matrix units used in construction of (4.5)-(4.8) are not defined:

$$P_{row}(k,n+1)=P_{col}(m+1,k)=P^{+}(k,0)=P_{+}(0,k)=0,\quad k=0,1,\dots.$$

When deriving (4.11) we apply formulae (2.21) to the matrix units (4.4)

$$E^{\Lambda(r,s)}_{row/col} \;=\; E^{\Lambda(r,s+1)}_{row}+E^{\Lambda(r+1,s)}_{col}+E^{\Lambda^{+}(r,s)}+E^{\Lambda_{+}(r,s)}\,, \tag{4.12}$$

and then use (2.14) in the form

$$\rho_R(E^{\lambda}_{\alpha})R_k \;=\; \omega^{-1}(\ell_k)\,\rho_R(E^{\lambda}_{\alpha,\pi_k(\alpha)})\;-\;\frac{q^{-\ell_k}}{(\ell_k)_q}\,\rho_R(E^{\lambda}_{\alpha})\,,\qquad \lambda\vdash(k+1),\qquad \ell_k\equiv\ell_k\{^{\lambda}_{\alpha}\} \tag{4.13}$$

in order to transform the terms $\rho_R(E^{\Lambda(r,s)}_{row/col})\,R_t$ in expressions for $P_{row/col}(k,i-k)$. Using calculations analogous to (3.24), one can show that the term with the non-diagonal matrix unit entering the right hand side of (4.13) does not contribute to (4.11).

Note that expressions (4.12) for $E^{\Lambda(r,s)}_{row}$ and $E^{\Lambda(r,s)}_{col}$ are *formally* the same, since our notations (4.4) take into account only the position of the largest integer among those filling the Young tableau. One should keep in mind that the matrix units $E^{\Lambda(r,s)}_{row}$ and $E^{\Lambda(r,s)}_{col}$ in the right hand side of (4.12) differ in the position of the number preceding to the largest one. This difference manifests itself in applying relation (4.13).

At the next step we reduce the similar terms in (4.11):

$$\begin{aligned}
\Phi_i &= -\sum_{k=\max\{0,i-n\}}^{\min\{i,m\}} (-1)^k\, q^{2k-i}\,\{P^+(k,i-k)+P_+(k,i-k)\} \\
&+ \sum_{k=\max\{0,i+1-n\}}^{\min\{i,m\}} \frac{(-1)^k}{(m+n-i+2)_q}\Big\{q^{-(m+n-i+1)}\,\frac{k_q(m+n-k+2)_q}{(m+n-i+1)_q} \\
&\qquad + q\,(i-k)_q(m+n-i+k+2)_q\Big\}P_{row}(k,i-k+1) \\
&+ \sum_{k=\max\{0,i-n\}}^{\min\{i,m-1\}} \frac{(-1)^k}{(m+n-i+2)_q}\Big\{q^{(m+n-i+1)}\,\frac{(i-k)_q(m+n-i+k+2)_q}{(m+n-i+1)_q} \\
&\qquad + q^{-1}k_q(m+n-k+2)_q\Big\}P_{col}(k+1,i-k). \qquad (4.14)
\end{aligned}$$

In calculating the coefficients at $P^+(\cdot,\cdot)$ and $P_+(\cdot,\cdot)$ the following relation is useful

$$q^x y_q + q^{-y}x_q = (x+y)_q, \quad x,y\in\mathbb{Z}. \qquad (4.15)$$

Next, we transform the first summand in the right hand side of (4.14) with the help of the computation

$$\begin{aligned}
&P^+(k,i-k)+P_+(k,i-k)+P_{row}(k,i-k+1)+P_{col}(k+1,i-k) \\
&\quad = \mathrm{Tr}_{R^{(2\ldots A)}}\left(M_{\overline{2}}\ldots M_{\overline{A}}\;\rho_R(E^{\Lambda(r,s)}_{\alpha})\,R_{m+n-i}\ldots R_1\right) \qquad (4.16) \\
&\quad = \mathrm{Tr}_{R^{(2\ldots(m+n-i+1))}}\left(M_{\overline{2}}\ldots M_{\overline{m+n-i+1}}\,R_{m+n-i}\ldots R_1\right)\; s_{\Lambda(k,i-k)}(M) \\
&\quad = \phi(M^{\overline{m+n-i}})\,s_{\Lambda(k,i-k)}(M)\,. \qquad (4.17)
\end{aligned}$$

In the above computation we use relation (4.12) first what results in (4.16). Expression (4.16) turns out to be independent on a particular choice of the index α of the matrix unit. Then we split (4.16) into two factors with the use of (3.19). At last, the first factor is transformed in the same way as that used to derive relation (3.28) for the matrix powers from definition (3.27).

Substituting (4.17) into (4.14) one gets

$$\begin{aligned}\Phi_i &= -\phi(M^{\overline{m+n-i}})\sum_{k=max\{0,i-n\}}^{\min\{i,m\}}(-1)^k q^{2k-i}\, s_{\Lambda(k,i-k)}(M)\\ &+\sum_{k=\max\{0,i+1-n\}}^{\min\{i,m\}}(-1)^k\,\frac{(i+1-k)_q(m+n-i+k+1)_q}{(m+n-i+1)_q}\,P_{row}(k,i+1-k)\\ &+\sum_{k=\max\{0,i-n\}}^{\min\{i,m-1\}}(-1)^k\,\frac{(k+1)_q(m+n-k+1)_q}{(m+n-i+1)_q}\,P_{col}(k+1,i-k)\ . \qquad (4.18)\end{aligned}$$

Here to simplify the coefficients at $P_{row}(\cdot,\cdot)$ and $P_{col}(\cdot,\cdot)$ we have used the formula

$$q^{-x}\,\tfrac{(x+y+1)_q z_q}{x_q}+q\,y_q(x+z+1)_q+q^{z-y}\,(x+1)_q \;=\; \tfrac{(x+1)_q(y+1)_q(x+z)_q}{x_q}, \qquad (4.19)$$

which can be easily verified with the help of (4.15).

Finally, shifting the summation index $k\to k+1$ in the last summand of (4.18) we come to (4.10).

Now it remains only to construct analogs of (4.10) for the boundary values $i=0$ and $i=m+n$.

For $i=0$ we have:

$$\Phi_1=P_{row}(1,0)+P_{col}(0,1)=\phi(M^{\overline{m+n}})\,s_{\Lambda(0,0)}(M)\,, \qquad (4.20)$$

which coincides with (4.10) if one sets $\Phi_0=0$.

In case $i=m+n$ the transformation of Φ_{m+n} is performed in the same way as when deriving (4.10). A peculiarity of the case involved consists in two points. Firstly, when carrying out the computations one finds an additional term associated with the Young diagram $((n+1)^{m+1})$ and secondly, $\Phi_{m+n+1}=0$. So, making the calculations we find

$$\begin{aligned}\Phi_{m+n} &= (-1)^{m+1}q^{m-n}\,\mathrm{Id}_V\, s_{\Lambda(m,n)}(M)\\ &+ (-1)^m\,(m+1)_q(n+1)_q\mathrm{Tr}_{R^{(2\dots A)}}\Big\{M_{\overline{2}}\dots M_{\overline{A}}\,\rho_R\left(E_{m+1}^{((n+1)^{m+1})}\right)\Big\}, \qquad (4.21)\end{aligned}$$

which also has the same form as (4.10) provided that $\rho_R\left(E_{m+1}^{((n+1)^{m+1})}\right)=0$ or, by definition, in the $GL(m|n)$ case.

Next, we add together the left and right hand sides of all relations (4.10), (4.20) and then subtract from the sum the corresponding parts of (4.21). At last, on applying the map ϕ^{-1} (see (3.13), (3.14)) to the result we come to the following theorem.

Theorem 10 *(The Cayley-Hamilton identity) Let M be the matrix of generators of a quantum matrix algebra $\mathcal{M}(R,F)$ of the $GL(m|n)$ type. Then the following matrix identity holds*

$$\sum_{i=0}^{n+m} M^{\overline{m+n-i}} \sum_{k=\max\{0,i-n\}}^{\min\{i,m\}} (-1)^k\, q^{2k-i}\, s_{\Lambda(k,i-k)}(M) \equiv 0\,, \tag{4.22}$$

If one multiplies the both sides of the matrix equalities (4.10), (4.20) and (4.21) by the matrix M from the left, the equalities turn into relations among the elements of the space $\mathrm{Pow}(R,F)$. Adding this relations together as in deriving the Cayley-Hamilton theorem, we establish the connection among the elements of two basis sets of the space $\mathrm{Pow}(R,F)$ which were described in proposition 9.

Proposition 11 *Let M be the matrix of generators of a quantum matrix algebra $\mathcal{M}(R,F)$ of the Hecke type. Then the λ-powers of M corresponding to the rectangular Young diagrams $\lambda = ((r+1)^{s+1})$, $r,s = 0,1,\ldots$ are expressed in terms of its $\overline{k}$-th powers as follows*

$$\begin{aligned}(-1)^s\,(s+1)_q (r+1)_q M^{\left(((r+1)^{s+1});\, s+1\right)} \\ = \sum_{i=0}^{s+r} M^{\overline{s+r+1-i}} \sum_{k=\max\{0,i-r\}}^{\min\{i,s\}} (-1)^k\, q^{2k-i}\, s_{\Lambda(k,i-k)}(M).\end{aligned} \tag{4.23}$$

References

[DKS] Damaskinsky E.V., Kulish P.P., Sokolov M.A., *'Gauss Decomposition for Quantum Groups and Supergroups'* Zap. Nauch. Semin. POMI **211** (1995) 11–45.

[DJ] Dipper R. and James G., *'Representations of Hecke algebras of general linear groups'*, Proc. London Math. Soc. (3) **52** (1986) 20-52.
Dipper R. and James G., *'Blocks and idempotents of Hecke algebras of general linear groups'*, Proc. London Math. Soc. (3) **54** (1987) 57-82.

[D] Drinfeld V.G., *'Quasi-Hopf Algebras'*. Leningrad Math. J. **1** (1990) 1419–1457.

[FP] Faddeev L.D., Pyatov P.N., *'The Differential Calculus on Quantum Linear Groups'* Trans. Amer. Math. Soc. Ser. 2 **175** (1996) 35–47.

[FRT] Reshetikhin N.Yu., Takhtajan L.A, Faddeev L.D., *'Quantization of Lie groups and Lie algebras'*, Algebra i Analiz, **1**, no. 1 (1989) 178–206 (in Russian).

[Gou] Gould M.D., *'On the matrix elements of the $U(n)$ generators'*. J. Math. Phys. **22** (1981) 15–22.

[G] Gurevich D.I., *'Algebraic aspects of the Yang-Baxter equation'*, Algebra i Analiz, **62**, no. 4 (1990) 119 – 148 (in Russian).

[GPS] Gurevich D., Pyatov P., Saponov P., *'Hecke symmetries and characteristic relations on reflection equation algebras'*. Lett. Math. Phys. **41** (1997) 255–264.

[GS] Gurevich D., Saponov P., *'Quantum line bundles via Cayley-Hamilton identity'*, J. Phys. A: Math. Gen. **34** no. 21 (2001) 4553 – 4569.

[GS2] Gurevich D., Saponov P., *'Quantum line bundles on a noncommutative sphere'* J. Phys. A: Math. Gen. **35** (2002) 9629–9643.

[I] Isaev A.P., *'Quantum groups and Yang-Baxter equations,'* Sov. J. Part. Nucl. **26** (1995) 501–526.

[IOP1] Isaev A.P., Ogievetsky O.V., Pyatov P.N., *'On quantum matrix algebras satisfying the Cayley-Hamilton-Newton identities'*. J. Phys. A: Math. Gen. **32** (1999) L115-L121.

[IOP2] Isaev A.P., Ogievetsky O.V., Pyatov P.N., *'Cayley-Hamilton-Newton Identities and Quasitriangular Hopf Algebras'*. In Proc. of International Workshop "Supersymmetries and Quantum Symmetries", July 27-31, 1999. Eds. E.Ivanov, S.Krivonos and A.Pashnev, JINR, Dubna E2-2000-82, pp. 397–405; Preprint `math.QA/9912197`.

[IOPS] Isaev A.P., Ogievetsky O.V., Pyatov P.N., Saponov P.A., *'Characteristic Polynomials for Quantum Matrices'*. In Proc. of International Conference in memory of V.I.Ogievetsky 'Supersymmetries and Quantum symmetries', (Dubna, Russia, 1997). Eds. J. Wess and E. Ivanov, Lecture Notes in Physics, vol. **524**, pp. 322 – 330, Springer Verlag, 1998.

[IP] Isaev A.P., Pyatov P.N., *'Covariant differential complexes on quantum linear groups'* J. Phys. A: Math. Gen. **28** (1995) 2227-2246.

[KT] Kantor I., Trishin I., *'On a concept of determinant in the supercase'*. Comm. in Algebra, **22** (1994) 3679 – 3739.
Kantor I., Trishin I., *'On the Cayley-Hamilton equation in the supercase'*. Comm. in Algebra, **27** (1999) 233 – 259.

[KS] Kulish P.P., Sasaki R., *'Covariance Properties of Reflection Equation Algebras'*. Prog. Theor. Phys., **89** (1993) 741 – 761.

[Mac] Macdonald I. G., *'Symmetric Functions and Hall Polynomials (Oxford Mathematical Monographs)'*, Oxford University Press, 1998.

[M] Murphy G. E., *'On the representation theory of the symmetric groups and associated Hecke algebras'*, J. Algebra **152** (1992) 492-513.
Murphy G. E., *'The representations of Hecke algebras of Type A_n'*, J. Algebra **173** (1995) 97-121.

[NT] Nazarov M., Tarasov V., *'Yangians and Gelfand-Zetlin bases'*. Publ. RIMS **30** (1994) 459.

[OP] Ogievetsky O. and Pyatov P., *'Lecture on Hecke algebras'*, in Proc. of the International School "Symmetries and Integrable Systems" Dubna, Russia, June 8-11, 1999. JINR, Dubna, D2,5-2000-218, pp.39-88; Preprint CPT-2000/P.4076 and MPI 01-40.

[OP2] Ogievetsky O.V., Pyatov P.N.,*'Cayley-Hamilton identities for quantum matrix algebras of Birman-Murakami-Wenzl type'*, to appear.

[PS] Pyatov P., Saponov P., *'Characteristic relations for quantum matrices'*. J. Phys. A: Math. Gen. **28** (1995) 4415 – 4421.

[R] Ram A., *'Seminormal representations of Weyl groups and Iwahori-Hecke algebras'*, Proc. of London Math. Soc. **75** (1997) 99-133, [arXiv:math.RT/9511223].